

Present in more than 30 countries,
a worldwide reference
for honey bee health and nutrition



Chemicals Laif

Viale dell'Artigianato, 13 - 35010 Vigonza (PD) - Italia



✉ info@chemicalslaif.it

🌐 www.alveis.it

☎ +39 049 626281

ApilifeVar®




VETERINARY MEDICINE BASED ON ESSENTIAL OILS AGAINST VARROA

NO
DRUG
RESISTANCE

NO
TOXIC
COMPOUNDS

SUITABLE
FOR ANY
TYPE OF
BEEHIVE



API LIFE VAR®
bee-hive strip for honey bees

Each bee-hive strip contains:
Active substances: Thymol 8,00 g; Eucalyptus oil 1,72g; Camphor racemic 0,39 g; Levomenthol 0,39g;
Excipients: foamed phenolic resin.

Pharmaceutical form: bee-hive strip.
Packaging size: sachet containing 2 strips
Target species: Bees (Apis mellifera)

Indications: Treatment of varroosis of the bees caused by Varroa destructor.

Dosage, route and method of administration:
Dosage: 1 strip every 7 days for each beehive, complete treatment 4 strips for each beehive. The treatment should be done once per year.

Application: to perform the treatment take one strip from the sachet and place it in one of the four corners above the honeycombs, far away from the brood placed in the center of the beehive. Close the beehive and let the product work for 7 days. The strip can also be used by cutting it into 3-4 pieces to be placed in the corners of the beehive. Repeat the treatment for 4 consecutive times. At the end of the treatment, the pieces of strips still present must be removed. It is not recommended to use the product in multiple storey hives as unsatisfactory efficacy might be expected.

Withdrawal period: Honey: Zero days
Do not use during honey flow in order to avoid adulterate the taste of the honey.

Adverse reactions: Bees may remove food from directly under the strip. Treatment at temperatures above 30°C may lead to increased stress and mortality of adult bees and brood. The colony may become slightly agitated during treatment. There may be reduced acceptance of the food if feeding at the same time as treatment. If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

Contraindications: Do not use during honey flow in order to avoid tainting the taste of the honey. Do not use when maximum daytime temperature is above 30°C.


FOR ANIMAL TREATMENT ONLY

AVM-GSL

Marketing authorisation holder:
CHEMICALS LAIF s.r.l.
V.le dell'Artigianato n° 13, 35010 Vigonza (PD),
Tel. +39 049 626281 - Fax +39 049 628501
e-mail: info@chemicalslaif.it

Marketing authorisation number: Vm 23101

Batch No - Expiry date: see on the back of the sachet.
After first opening of the sachet all strips should be used immediately.



APRI QUI (OPEN HERE)



NO
drug
resistance

NO
toxic
compounds

>94%
efficacy
against
Varroa

ApiLifeVar®

**WHEN
TO USE IT**

SPRING - SUMMER

**IDEAL
TEMPERATURE
20-25°C**
Range 15-30° C

**HOW
TO USE IT**

1 STRIPS FOR BEEHIVE

weekly for
4
WEEKS

ADVANTAGES:


- no need for extra space in the hive
- dosage suitable to any type of hive
- easy and safe to use
- excellent efficacy and tolerability
- allowed in organic beekeeping

**THE ONLY
VETERINARY MEDICINE WITH
THE SYNERGIC ACTION OF
4 ACTIVE INGREDIENTS**

ApiLife Var is the **organic
solution against varroa.**
Adaptable to all climate
conditions, kind of bee
breeding and beehives.

APILIFE VAR Veterinary medicine

based on essential oils
(Thymol, Eucalyptus oil, Levomenthol, Racemic camphor)



1 sachet = 2 strips


WAY OF USE:



1
Open the bag
containing the
two strips.



2
Take one of the two
strips and break it in
2-4 parts.



3
Open the beehive and set the
single pieces of the strip in the
corners of the beehive.

4

Close the beehive and let the
product work for 7 days.
Repeat the treatment for 4
times with the other strips.


ADVICE FOR USE:

- Balance the beehives according to their strenght before the use
- All colonies in the same apiary should be treated simultaneously to avoid robbery
- Treat by introducing the product into the beehive either early in the morning or late in the afternoon towards sunset, avoiding the hottest hours
- Do the treatment without supers
- Do not place the product in the center of the beehive or near the brood.
- Handle with care the package

For further instructions and advice of use see the label of the product

APILIFE VAR FIELD TESTS

Effectiveness against varroa in different countries



Experience over many years using the product as well as numerous studies carried out in Italy and internationally prove the high effectiveness of ApiLife Var® showing an average Varroa mite drop of higher than 94%.

